

# Some identities of symmetry for the generalized Bernoulli numbers and polynomials

By

Taekyun KIM

**Abstract.** In this paper, by the properties of  $p$ -adic invariant integral on  $\mathbb{Z}_p$ , we establish various identities concerning the generalized Bernoulli numbers and polynomials. From the symmetric properties of  $p$ -adic invariant integral on  $\mathbb{Z}_p$ , we give some interesting relationship between the power sums and the generalized Bernoulli polynomials.

**2000 Mathematics Subject Classification:** 11B68, 11M38, 11S80.

**Key Words and Phrases:**  $p$ -adic invariant integral, Bernoulli numbers, Bernoulli polynomials.

## §1. Introduction

Let  $p$  be a fixed prime number. Throughout this paper, the symbols  $\mathbb{Z}$ ,  $\mathbb{Z}_p$ ,  $\mathbb{Q}_p$ , and  $\mathbb{C}_p$  will denote the ring of rational integers, the ring of  $p$ -adic integers, the field of  $p$ -adic rational numbers, and the completion of algebraic closure of  $\mathbb{Q}_p$ , respectively. Let  $\mathbb{N}$  be the set of natural numbers and  $\mathbb{Z}_+ = \mathbb{N} \cup \{0\}$ . Let  $v_p$  be the normalized exponential valuation of  $\mathbb{C}_p$  with  $|p|_p = p^{-v_p(p)} = 1/p$ . Let  $UD(\mathbb{Z}_p)$  be the space of uniformly differentiable function on  $\mathbb{Z}_p$ . For  $f \in UD(\mathbb{Z}_p)$ , the  $p$ -adic invariant integral on  $\mathbb{Z}_p$  is defined as

$$I(f) = \int_{\mathbb{Z}_p} f(x) dx = \lim_{N \rightarrow \infty} \frac{1}{p^N} \sum_{x=0}^{p^N-1} f(x), \quad (\text{see [6]}). \quad (1)$$

From the definition (1), we have

$$I_1(f_1) = I_1(f) + f'(0), \text{ where } f'(0) = \frac{df(x)}{dx}|_{x=0} \text{ and } f_1(x) = f(x+1). \quad (2)$$

Let  $f_n(x) = f(x+n)$ , ( $n \in \mathbb{N}$ ). Then we can derive the following equation (3) from (2).

$$I(f_n) = I(f) + \sum_{i=0}^n f'(i), \quad (\text{see [6]}). \quad (3)$$

It is well known that the ordinary Bernoulli polynomials  $B_n(x)$  are defined as

$$\frac{t}{e^t - 1} e^{xt} = \sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!}, \quad (\text{ see [1-25] } ),$$

and the Bernoulli number  $B_n$  are defined as  $B_n = B_n(0)$ .

Let  $d$  a fixed positive integer. For  $n \in \mathbb{N}$ , we set

$$X = X_d = \varprojlim_{\mathbb{N}} \left( \mathbb{Z}/dp^N \mathbb{Z} \right), \quad X_1 = \mathbb{Z}_p;$$

$$X^* = \bigcup_{\substack{0 < a < dp, \\ (a,p)=1}} (a + dp\mathbb{Z}_p);$$

$$a + dp^N \mathbb{Z}_p = \{ x \in X \mid x \equiv a \pmod{dp^N} \},$$

where  $a \in \mathbb{Z}$  lies in  $0 \leq a < dp^N$ . In [6], it is known that

$$\int_X f(x) dx = \int_{\mathbb{Z}_p} f(x) dx, \quad \text{for } f \in UD(\mathbb{Z}_p).$$

Let us take  $f(x) = e^{tx}$ . Then we have

$$\int_{\mathbb{Z}_p} e^{tx} dx = \frac{t}{e^t - 1} = \sum_{n=0}^{\infty} B_n \frac{t^n}{n!}.$$

Thus, we note that

$$\int_{\mathbb{Z}_p} x^n dx = B_n, \quad n \in \mathbb{Z}_+, \quad (\text{see [1-25]}).$$

Let  $\chi$  be the Dirichlet's character with conductor  $d \in \mathbb{N}$ . Then the generalized Bernoulli polynomials attached to  $\chi$  are defined as

$$\sum_{a=1}^d \frac{\chi(a) t e^{at}}{e^{dt} - 1} e^{xt} = \sum_{n=0}^{\infty} B_{n,\chi}(x) \frac{t^n}{n!}, \quad (\text{ see [22] } ), \quad (4)$$

and the generalized Bernoulli numbers attached to  $\chi$ ,  $B_{n,\chi}$  are defined as  $B_{n,\chi} = B_{n,\chi}(0)$ .

In this paper, we investigate the interesting identities of symmetry for the generalized Bernoulli numbers and polynomials attached to  $\chi$  by using the properties of  $p$ -adic invariant integral on  $\mathbb{Z}_p$ . Finally, we will give relationship between the power sum polynomials and the generalized Bernoulli numbers attached to  $\chi$ .

## §2. Symmetry of power sum and the generalized Bernoulli polynomials

Let  $\chi$  be the Dirichlet character with conductor  $d \in \mathbb{N}$ . From (3), we note that

$$\int_X \chi(x) e^{xt} dx = \frac{t \sum_{i=0}^{d-1} \chi(i) e^{it}}{e^{dt} - 1} = \sum_{n=0}^{\infty} B_{n,\chi} \frac{t^n}{n!}, \quad (5)$$

where  $B_{n,\chi}(x)$  are  $n$ -th generalized Bernoulli numbers attached to  $\chi$ . Now, we also see that the generalized Bernoulli polynomials attached to  $\chi$  are given by

$$\int_X \chi(y) e^{(x+y)t} dy = \frac{\sum_{i=0}^{d-1} \chi(i) e^{it}}{e^{dt} - 1} e^{xt} = \sum_{n=0}^{\infty} B_{n,\chi}(x) \frac{t^n}{n!}. \quad (6)$$

By (5) and (6), we easily see that

$$\int_X \chi(x) x^n dx = B_{n,\chi}, \quad \text{and} \quad \int_X \chi(y) (x+y)^n dy = B_{n,\chi}(x). \quad (7)$$

From (6), we have

$$B_{n,\chi}(x) = \sum_{\ell=0}^n \binom{n}{\ell} B_{\ell,\chi} x^{n-\ell}. \quad (8)$$

From (6), we can also derive

$$\int_X \chi(x) e^{xt} dx = \sum_{i=0}^{d-1} \chi(i) \frac{t}{e^{dt} - 1} e^{(\frac{i}{d})dt} = \sum_{n=0}^{\infty} \left( d^n \sum_{i=0}^{d-1} \chi(i) B_n\left(\frac{i}{d}\right) \right) \frac{t^n}{n!}.$$

Therefore, we obtain the following lemma.

LEMMA1. *For  $n \in \mathbb{Z}_+$ , we have*

$$\int_X \chi(x) x^n dx = B_{n,\chi} = d^n \sum_{i=0}^{d-1} \chi(i) B_i\left(\frac{i}{d}\right).$$

We observe that

$$\frac{1}{t} \left( \int_X \chi(x) e^{(nd+x)t} dx - \int_X e^{xt} \chi(x) dx \right) = \frac{nd \int_X \chi(x) e^{xt} dx}{\int_X e^{ndxt} dx} = \frac{e^{ndt} - 1}{e^{dt} - 1} \left( \sum_{i=0}^{d-1} \chi(i) e^{it} \right). \quad (9)$$

Thus, we have

$$\frac{1}{t} \left( \int_X \chi(x) e^{(nd+x)t} dx - \int_X e^{xt} dx \right) = \sum_{k=0}^{\infty} \left( \sum_{\ell=0}^{nd-1} \chi(\ell) \ell^k \right) \frac{t^k}{k!}. \quad (10)$$

Let us define the  $p$ -adic functional  $T_k(\chi, n)$  as follows:

$$T_k(\chi, n) = \sum_{\ell=0}^n \chi(\ell) \ell^k, \quad \text{for } k \in \mathbb{Z}_+. \quad (11)$$

By (10) and (11), we see that

$$\frac{1}{t} \left( \int_X \chi(x) e^{(nd+x)t} dx - \int_X e^{xt} dx \right) = \sum_{n=0}^{\infty} \left( T_k(\chi, nd-1) \right) \frac{t^k}{k!}. \quad (12)$$

By using Taylor expansion in (12), we have

$$\int_X \chi(x) (dn+x)^k dx - \int_X \chi(x) x^k dx = k T_{k-1}(\chi, nd-1), \quad \text{for } k, n, d \in \mathbb{N}. \quad (13)$$

That is,

$$B_{k,\chi}(nd) - B_{k,\chi} = k T_{k-1}(\chi, nd-1).$$

Let  $w_1, w_2, d \in \mathbb{N}$ . Then we consider the following integral equation

$$\begin{aligned} & \frac{d \int_X \int_X \chi(x_1) \chi(x_2) e^{(w_1 x_1 + w_2 x_2)t} dx_1 dx_2}{\int_X e^{dw_1 w_2 x t} dx} \\ &= \frac{t(e^{dw_1 w_2 t} - 1)}{(e^{w_1 d t} - 1)(e^{w_2 d t} - 1)} \left( \sum_{a=0}^{d-1} \chi(a) e^{w_1 a t} \right) \left( \sum_{b=0}^{d-1} \chi(b) e^{w_2 b t} \right). \end{aligned} \quad (14)$$

From (9) and (12), we note that

$$\frac{dw_1 \int_X \chi(x) e^{xt} dx}{\int_X e^{dw_1 x t} dx} = \sum_{k=0}^{\infty} \left( T_k(\chi, dw_1 - 1) \right) \frac{t^k}{k!}. \quad (15)$$

Let us consider the  $p$ -adic functional  $T_\chi(w_1, w_2)$  as follows:

$$T_\chi(w_1, w_2) = \frac{d \int_X \int_X \chi(x_1) \chi(x_2) e^{(w_1 x_1 + w_2 x_2 + w_1 w_2 x)t} dx_1 dx_2}{\int_X e^{dw_1 w_2 x_3 t} dx_3}. \quad (16)$$

Then we see that  $T_\chi(w_1, w_2)$  is symmetric in  $w_1$  and  $w_2$ , and

$$T_\chi(w_1, w_2) = \frac{t(e^{dw_1 w_2 t} - 1) e^{w_1 w_2 x t}}{(e^{w_1 d t} - 1)(e^{w_2 d t} - 1)} \left( \sum_{a=0}^{d-1} \chi(a) e^{w_1 a t} \right) \left( \sum_{b=0}^{d-1} \chi(b) e^{w_2 b t} \right). \quad (17)$$

By (16) and (17), we have

$$\begin{aligned}
T_\chi(w_1, w_2) &= \left( \frac{1}{w_1} \int_X \chi(x_1) e^{w_1(x_1+w_2x)t} dx_1 \right) \left( \frac{dw_1 \int_X \chi(x_2) e^{w_2x_2t} dx_2}{\int_X e^{dw_1w_2xt} dx} \right) \\
&= \left( \frac{1}{w_1} \sum_{i=0}^{\infty} B_{i,\chi}(w_2x) \frac{w_1^i t^i}{i!} \right) \left( \sum_{k=0}^{\infty} T_k(\chi, dw_1 - 1) \frac{w_2^k t^k}{k!} \right) \\
&= \frac{1}{w_1} \left( \sum_{\ell=0}^{\infty} \left( \sum_{i=0}^{\ell} \frac{B_{i,\chi}(w_2x) T_{\ell-i}(\chi, dw_1 - 1) w_1^i w_2^{\ell-i} \ell!}{i! (\ell-i)!} \right) \frac{t^\ell}{\ell!} \right) \\
&= \sum_{\ell=0}^{\infty} \left( \sum_{i=0}^{\ell} \binom{\ell}{i} B_{i,\chi}(w_2x) T_{\ell-i}(\chi, dw_1 - 1) w_1^{i-1} w_2^{\ell-i} \right) \frac{t^\ell}{\ell!}.
\end{aligned} \tag{18}$$

From the symmetric property of  $T_\chi(w_1, w_2)$  in  $w_1$  and  $w_2$ , we note that

$$\begin{aligned}
T_\chi(w_1, w_2) &= \left( \frac{1}{w_2} \int_X \chi(x_2) e^{w_2(x_2+w_1x)t} dx_2 \right) \left( \frac{dw_2 \int_X \chi(x_1) e^{w_1x_1t} dx_1}{\int_X e^{dw_1w_2xt} dx} \right) \\
&= \left( \frac{1}{w_2} \sum_{i=0}^{\infty} B_{i,\chi}(w_1x) \frac{w_2^i t^i}{i!} \right) \left( \sum_{k=0}^{\infty} T_k(\chi, dw_2 - 1) \frac{w_1^k t^k}{k!} \right) \\
&= \frac{1}{w_2} \left( \sum_{\ell=0}^{\infty} \left( \sum_{i=0}^{\ell} \frac{B_{i,\chi}(w_1x) w_2^i T_{\ell-i}(\chi, dw_2 - 1) w_1^{\ell-i} \ell!}{i! (\ell-i)!} \right) \frac{t^\ell}{\ell!} \right) \\
&= \sum_{\ell=0}^{\infty} \left( \sum_{i=0}^{\ell} \binom{\ell}{i} w_2^{i-1} w_1^{\ell-i} B_{i,\chi}(w_1x) T_{\ell-i}(\chi, dw_2 - 1) \right) \frac{t^\ell}{\ell!}.
\end{aligned} \tag{19}$$

By comparing the coefficients on the both sides of (18) and (19), we obtain the following theorem.

**THEOREM 2.** *For  $w_1, w_2, d \in \mathbb{N}$ , we have*

$$\begin{aligned}
&\sum_{i=0}^{\ell} \binom{\ell}{i} B_{i,\chi}(w_2x) T_{\ell-i}(\chi, dw_1 - 1) w_1^{i-1} w_2^{\ell-i} \\
&= \sum_{i=0}^{\ell} \binom{\ell}{i} B_{i,\chi}(w_1x) T_{\ell-i}(\chi, dw_2 - 1) w_2^{i-1} w_1^{\ell-i}.
\end{aligned}$$

Let  $x = 0$  in Theorem 2. Then we have

$$\begin{aligned}
&\sum_{i=0}^{\ell} \binom{\ell}{i} B_{i,\chi} T_{\ell-i}(\chi, dw_1 - 1) w_1^{i-1} w_2^{\ell-i} \\
&= \sum_{i=0}^{\ell} \binom{\ell}{i} B_{i,\chi} T_{\ell-i}(\chi, dw_2 - 1) w_2^{i-1} w_1^{\ell-i}.
\end{aligned}$$

By (15) and (17), we also see that

$$\begin{aligned}
T_\chi(w_1, w_2) &= \left( \frac{e^{w_1 w_2 x t}}{w_1} \int_X \chi(x_1) e^{w_1 x_1 t} dx_1 \right) \left( \frac{dw_1 \int_X \chi(x_2) e^{w_2 x_2 t} dx_2}{\int_X e^{dw_1 w_2 x t} dx} \right) \\
&= \left( \frac{e^{w_1 w_2 x t}}{w_1} \int_X \chi(x_1) e^{w_1 x_1 t} dx_1 \right) \left( \frac{e^{dw_1 w_2 t} - 1}{e^{w_2 dt} - 1} \right) \left( \sum_{i=0}^{d-1} \chi(i) e^{w_2 i t} \right) \\
&= \left( \frac{e^{w_1 w_2 x t}}{w_1} \int_X \chi(x_1) e^{w_1 x_1 t} dx_1 \right) \left( \sum_{\ell=0}^{w_1-1} \sum_{i=0}^{d-1} e^{w_2(i+\ell d)t} \chi(i+\ell d) \right) \\
&= \left( \frac{e^{w_1 w_2 x t}}{w_1} \int_X \chi(x_1) e^{w_1 x_1 t} dx_1 \right) \left( \sum_{i=0}^{dw_1-1} e^{w_2 i t} \chi(i) \right) \tag{20} \\
&= \frac{1}{w_1} \sum_{i=0}^{dw_1-1} \chi(i) \int_X \chi(x_1) e^{w_1(x_1 + w_2 x + \frac{w_2}{w_1} i)t} dx_1 \\
&= \frac{1}{w_1} \sum_{i=0}^{dw_1-1} \chi(i) \sum_{k=0}^{\infty} B_{k,\chi}(w_2 x + \frac{w_2}{w_1} i) \frac{w_1^k t^k}{k!} \\
&= \sum_{k=0}^{\infty} \left( \sum_{i=0}^{dw_1-1} \chi(i) B_{k,\chi}(w_2 x + \frac{w_2}{w_1} i) w_1^{k-1} \right) \frac{t^k}{k!}.
\end{aligned}$$

From the symmetric property of  $T_\chi(w_1, w_2)$  in  $w_1$  and  $w_2$ , we can also derive the following equation.

$$\begin{aligned}
T_\chi(w_1, w_2) &= \left( \frac{e^{w_1 w_2 x t}}{w_2} \int_X \chi(x_2) e^{w_2 x_2 t} dx_2 \right) \left( \frac{dw_2 \int_X \chi(x_1) e^{w_1 x_1 t} dx_1}{\int_X e^{dw_1 w_2 x t} dx} \right) \\
&= \left( \frac{e^{w_1 w_2 x t}}{w_2} \int_X \chi(x_2) e^{w_2 x_2 t} dx_2 \right) \left( \frac{e^{dw_1 w_2 t} - 1}{e^{w_1 dt} - 1} \right) \left( \sum_{i=0}^{d-1} \chi(i) e^{w_1 i t} \right) \\
&= \left( \frac{e^{w_1 w_2 x t}}{w_2} \int_X \chi(x_2) e^{w_2 x_2 t} dx_2 \right) \left( \sum_{\ell=0}^{w_2-1} e^{w_1 \ell t} \right) \left( \sum_{i=0}^{d-1} \chi(i) e^{w_1 i t} \right) \\
&= \frac{1}{w_2} \sum_{i=0}^{dw_2-1} \chi(i) \int_X \chi(x_2) e^{w_2(x_2 + w_1 x + \frac{w_1}{w_2} i)t} dx_2 \tag{21} \\
&= \frac{1}{w_2} \sum_{i=0}^{dw_2-1} \chi(i) \sum_{k=0}^{\infty} B_{k,\chi}(w_1 x + \frac{w_1}{w_2} i) \frac{w_2^k t^k}{k!} \\
&= \sum_{k=0}^{\infty} \left\{ \sum_{i=0}^{dw_2-1} \chi(i) B_{k,\chi}(w_1 x + \frac{w_1}{w_2} i) w_2^{k-1} \right\} \frac{t^k}{k!}.
\end{aligned}$$

By comparing the coefficients on the both sides of (20) and (21), we obtain the following theorem.

THEOREM 3. For  $w_1, w_2, d \in \mathbb{N}$ , we have

$$\sum_{i=0}^{dw_1-1} \chi(i) B_{k,\chi}(w_2 x + \frac{w_2}{w_1} i) w_1^{k-1} = \sum_{i=0}^{dw_2-1} \chi(i) B_{k,\chi}(w_1 x + \frac{w_1}{w_2} i) w_2^{k-1}.$$

REMARK. Let  $x = 0$  in Theorem 3. Then we see that

$$\sum_{i=0}^{dw_1-1} \chi(i) B_{k,\chi}(\frac{w_2}{w_1} i) w_1^{k-1} = \sum_{i=0}^{dw_2-1} \chi(i) B_{k,\chi}(\frac{w_1}{w_2} i) w_2^{k-1}.$$

If we take  $w_2 = 1$ , then we have

$$\sum_{i=0}^{dw_1-1} \chi(i) B_{k,\chi}(\frac{i}{w_1}) w_1^{k-1} = \sum_{i=0}^{d-1} \chi(i) B_{k,\chi}(w_1 i).$$

## REFERENCES

- [1] L. C. Carlitz, *q-Bernoulli numbers and polynomials*, Duke Math. J. **15** (1948), 987-1000.
- [2] M. Cenkci, Y. Sisek, V. Kurt, *Further remarks on multiple p-adic q-L-function of two variables*, Adv. Stud. Contemp. Math. **14** (2007), 49-68.
- [3] M. Cenkci, M. Can, V. Kurt, *Multiple two-variable q-L-function and its behavior at s = 0*, Russ. J. Math. Phys. **15(4)** (2008), 447-459.
- [4] T. Ernst, *Example of a q-umbral calculus*, Adv. Stud. Contemp. Math. **16(1)** (2008), 1-22.
- [5] A. S. Hegazi, M. Mansour, *A note on q-Bernoulli numbers and polynomials*, J. Nonlinear Math. Phys. **13** (2006), 9-18.
- [6] T. Kim, *q-Volkenborn Integration*, Russ. J. Math. Phys. **9** (2002), 288-299.
- [7] T. Kim, *Non-archimedean q-integrals associated with multiple Changhee q-Bernoulli polynomials*, Russ. J. Math. Phys. **10** (2003), 91-98.
- [8] T. Kim, *Power series and asymptotic series associated with the q-analog of the two-variable p-adic L-function*, Russ. J. Math. Phys. **12(2)** (2005), 186-196.

- [9] T. Kim, *Multiple  $p$ -adic  $L$ -function* Russ. J. Math. Phys. **13**(2) (2006), 151-157.
- [10] T. Kim,  *$q$ -Euler numbers and polynomials associated with  $p$ -adic  $q$ -integral*, J. Non-linear Math. Phys. **14** (1) (2007), 15-27.
- [11] T. Kim, *A note on  $p$ -adic  $q$ -integral on  $\mathbb{Z}_p$* , Adv. Stud. Contemp. Math. **15** (2007), 133-138.
- [12] T. Kim,  *$q$ -Bernoulli numbers and polynomials associated with Gaussian binomial coefficients*, Russ. J. Math. Phys. **15** (1) (2008), 51-57.
- [13] T. Kim, *On the symmetry of the  $q$ -Bernoulli polynomials*, Abstr. Appl. Anal. **2008** (2008), Article ID914367, 7 pages.
- [14] T. Kim, *Symmetries  $p$ -adic invariant integral on  $\mathbb{Z}_p$  for Bernoulli and Euler polynomials*, J. Difference Equ. Appl. **14** (12) (2008), 1267-1277.
- [15] T. Kim, *Note on  $q$ -Genocchi numbers and polynomials*, Adv. Stud. Contemp. Math. **17** (1) (2008), 9-15.
- [16] T. Kim, *Symmetry of power sum polynomials and multivariate fermionic  $p$ -adic invariant integral on  $\mathbb{Z}_p$* , Russ. J. Math. Phys. **16** (1) (2009), 51-54.
- [17] Y. -H. Kim, W. Kim, L.-C. Jang, *On the  $q$ -extension of Apostol-Euler numbers and polynomials*, Abstr. Appl. Anal. **2008** (2008), Article ID296159, 10 pages.
- [18] B. A. Kupershmidt, *Reflection symmetries of  $q$ -Bernoulli polynomials*, J. Nonlinear Math. Phys. **12** (2005), 412-422.
- [19] H. Ozden, Y. Simsek, S. -H. Rim, I. N. Cangul *A note on  $p$ -adic  $q$ -Euler measure*, Adv. Stud. Contemp. Math. **14** (2007), 233-239.
- [20] K. H. Park, Y.-H. Kim *On some arithmetical properties of the Genocchi numbers and polynomials*, Advances in Difference Equations. <http://www.hindawi.com/journals/ade/aip.195049.html>.
- [21] M. Schork, *A representation of the  $q$ -fermionic commutation relations and the limit  $q = 1$* , Russ. J. Math. Phys. **12** (3) (2005), 394-399.
- [22] Y. Simsek, *Theorems on twisted  $L$ -function and twisted Bernoulli numbers*, Adv. Stud. Contemp. Math. **11** (2005), 205-218.
- [23] Y. Simsek, *On  $p$ -adic twisted  $q$ - $L$ -functions related to generalized twisted Bernoulli numbers*, Russ. J. Math. Phys. **13**(3) (2006), 340-348.



- [24] Y. Simsek, *Complete sums of  $(h, q)$ -extension of the Euler polynomials and numbers*, arXiv:0707.2849v1[math.NT].
- [25] Y.-H. Kim, K.-W. Hwang. *Symmetry of power sum and twisted Bernoulli polynomials*, Adv. Stud. Contemp. Math. **18 (2)** (2009), 105-113.

Taekyun KIM

Division of General Education-Mathematics,  
Kwangwoon University, Seoul 139-701, S. Korea

*E-mail:* tkkim@kw.ac.kr